

# Package: EMpeaksR (via r-universe)

October 31, 2024

**Type** Package

**Title** Conducting the Peak Fitting Based on the EM Algorithm

**Version** 0.3.1

**Description** The peak fitting of spectral data is performed by using the frame work of EM algorithm. We adapted the EM algorithm for the peak fitting of spectral data set by considering the weight of the intensity corresponding to the measurement energy steps (Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2019, 2021 and 2023) <[doi:10.1080/14686996.2019.1620123](https://doi.org/10.1080/14686996.2019.1620123)>, <[doi:10.1080/27660400.2021.1899449](https://doi.org/10.1080/27660400.2021.1899449)>, <[doi:10.1080/27660400.2022.2159753](https://doi.org/10.1080/27660400.2022.2159753)>). The package efficiently estimates the parameters of Gaussian mixture model during iterative calculation between E-step and M-step, and the parameters are converged to a local optimal solution. This package can support the investigation of peak shift with two advantages: (1) a large amount of data can be processed at high speed; and (2) stable and automatic calculation can be easily performed.

**License** MIT + file LICENSE

**Language** en-US

**Encoding** UTF-8

**RoxygenNote** 7.1.1

**Suggests** testthat (>= 3.0.0)

**Config/testthat/edition** 3

**NeedsCompilation** no

**Author** Tarojiro Matsumura [aut, cre]

**Maintainer** Tarojiro Matsumura <[matsumura-tarojiro@aist.go.jp](mailto:matsumura-tarojiro@aist.go.jp)>

**Date/Publication** 2023-03-29 13:40:02 UTC

**Repository** <https://matsumuratarojiro.r-universe.dev>

**RemoteUrl** <https://github.com/cran/EMpeaksR>

**RemoteRef** HEAD

**RemoteSha** ffb248b9e25ef160ca41969bdd5df74e881b740c

## Contents

show_dsgmm_curve . . . . .	2
show_gmm_curve . . . . .	5
show_lmm_curve . . . . .	6
show_pvmm_curve . . . . .	8
show_pvmm_lback_curve . . . . .	10
spect_em_dsgmm . . . . .	12
spect_em_gmm . . . . .	15
spect_em_lmm . . . . .	17
spect_em_pvmm . . . . .	19
spect_em_pvmm_lback . . . . .	21

<b>Index</b>	<b>25</b>
--------------	-----------

---

show_dsgmm_curve	<i>Visualization of the result of spect_em_dsgmm</i>
------------------	--

---

## Description

Visualization of the result of spect\_em\_dsgmm().

## Usage

```
show_dsgmm_curve(spect_em_dsgmm_res,
                 x,
                 y,
                 mix_ratio_init,
                 mu_init,
                 sigma_init,
                 alpha_init,
                 eta_init)
```

## Arguments

spect_em_dsgmm_res	data set obtained by spect_em_dsgmm()
x	measurement steps
y	intensity
mix_ratio_init	initial values of the mixture ratio of the components
mu_init	initial values of the mean of the components
sigma_init	initial values of the standard deviation of the components
alpha_init	initial values of the asymmetric parameter of the components
eta_init	initial values of the mixing ratio of Gauss and Lorentz distribution

**Details**

Perform a visualization of fitting curve estimated by Doniach-Sunjic-Gauss mixture model.

**Value**

Show the fitting curve and variation of the parameters.

**References**

Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2019). Spectrum adapted expectation-maximization algorithm for high-throughput peak shift analysis. *Science and technology of advanced materials*, 20(1), 733-745.

Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2021). Spectrum adapted expectation-conditional maximization algorithm for extending high-throughput peak separation method in XPS analysis. *Science and Technology of Advanced Materials: Methods*, 1(1), 45-55.

**Examples**

```
#generating the synthetic spectral data based on three component Doniach-Sunjic-Gauss mixture model.
x <- seq(0, 100, by = 0.5)
true_mu <- c(20, 50, 80)
true_sigma <- c(3, 3, 3)
true_alpha <- c(0.1, 0.3, 0.1)
true_eta <- c(0.4, 0.6, 0.1)
true_mix_ratio <- rep(1/3, 3)
degree <- 4

#truncated Doniach-Sunjic-Gauss
truncated_dsg <- function(x, mu, sigma, alpha, eta) {
  ((eta*((gamma(1-alpha)) /
  ((x-mu)^2+(sqrt(2*log(2))*sigma)^2)^((1-alpha)/2)) *
  cos((pi*alpha/2)+(1-alpha)*atan((x-mu) /
  (sqrt(2*log(2))*sigma)))) + (1-eta)*dnorm(x, mu, sigma)) /
  sum( ((eta*((gamma(1-alpha)) /
  ((x-mu)^2+(sqrt(2*log(2))*sigma)^2)^((1-alpha)/2)) *
  cos((pi*alpha/2)+(1-alpha)*atan((x-mu) /
  (sqrt(2*log(2))*sigma)))) + (1-eta)*dnorm(x, mu, sigma))
}

y <- c(true_mix_ratio[1]*truncated_dsg(x = x,
  mu = true_mu[1],
  sigma = true_sigma[1],
  alpha = true_alpha[1],
  eta = true_eta[1])*10^degree +
  true_mix_ratio[2]*truncated_dsg(x = x,
  mu = true_mu[2],
  sigma = true_sigma[2],
  alpha = true_alpha[2],
  eta = true_eta[2])*10^degree +
  true_mix_ratio[3]*truncated_dsg(x = x,
  mu = true_mu[3],
```

```

sigma = true_sigma[3],
alpha = true_alpha[3],
eta = true_eta[3])*10^degree)

plot(y~x, main = "genrated synthetic spectral data")

#Peak fitting by EMpeaksR
#Initial values
K <- 3
mix_ratio_init <- c(0.2, 0.4, 0.4)
mu_init <- c(20, 40, 70)
sigma_init <- c(4, 3, 2)
alpha_init <- c(0.3, 0.2, 0.4)
eta_init <- c(0.5, 0.4, 0.3)

#Conducting calculation
SP_ECM_DSG_res <- spect_em_dsgmm(x = x,
                                y = y,
                                mu = mu_init,
                                sigma = sigma_init,
                                alpha = alpha_init,
                                eta = eta_init,
                                mix_ratio = mix_ratio_init,
                                conv.cri = 1e-2,
                                maxit = 2000)

#Plot fitting curve and trace plot of parameters
show_dsgmm_curve(SP_ECM_DSG_res,
                 x,
                 y,
                 mix_ratio_init,
                 mu_init,
                 sigma_init,
                 alpha_init,
                 eta_init)

#Showing the result of spect_em_dsgmm()
print(cbind(c(mu_init),
            c(sigma_init),
            c(alpha_init),
            c(eta_init),
            c(mix_ratio_init)))

print(cbind(SP_ECM_DSG_res$mu,
            SP_ECM_DSG_res$sigma,
            SP_ECM_DSG_res$alpha,
            SP_ECM_DSG_res$eta,
            SP_ECM_DSG_res$mix_ratio))

print(cbind(true_mu,
            true_sigma,
            true_alpha,
            true_eta,

```

```
true_mix_ratio))
```

---

show_gmm_curve	<i>Visualization of the result of spect_em_gmm</i>
----------------	--

---

## Description

Visualization of the result of spect\_em\_gmm().

## Usage

```
show_gmm_curve(spect_em_gmm_res, x, y, mix_ratio_init, mu_init, sigma_init)
```

## Arguments

spect_em_gmm_res	data set obtained by spect_em_gmm()
x	measurement steps
y	intensity
mix_ratio_init	initial values of the mixture ratio of the components
mu_init	initial values of the mean of the components
sigma_init	initial values of the standard deviation of the components

## Details

Perform a visualization of fitting curve estimated by Gaussian mixture model.

## Value

Show the fitting curve and variation of the parameters.

## References

Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2019). Spectrum adapted expectation-maximization algorithm for high-throughput peak shift analysis. *Science and technology of advanced materials*, 20(1), 733-745.

## Examples

```
#generating the synthetic spectral data based on three component Gaussian mixture model.
x <- seq(0, 100, by = 0.5)
true_mu <- c(35, 50, 65)
true_sigma <- c(3, 3, 3)
true_mix_ratio <- rep(1/3, 3)
degree <- 4

y <- c(true_mix_ratio[1] * dnorm(x = x, mean = true_mu[1], sd = true_sigma[1])*10^degree +
```

```

true_mix_ratio[2] * dnorm(x = x, mean = true_mu[2], sd = true_sigma[2])*10^degree +
true_mix_ratio[3] * dnorm(x = x, mean = true_mu[3], sd = true_sigma[3])*10^degree)

plot(y~x, main = "generated synthetic spectral data")

#Peak fitting by EMpeaksR
#Initial values
K <- 3

mix_ratio_init <- c(0.2, 0.4, 0.4)
mu_init <- c(20, 40, 70)
sigma_init <- c(2, 5, 4)

#Conducting calculation
SP_EM_G_res <- spect_em_gmm(x, y, mu = mu_init, sigma = sigma_init, mix_ratio = mix_ratio_init,
                           conv.cri = 1e-2, maxit = 2000)

#Plot fitting curve and trace plot of parameters
show_gmm_curve(SP_EM_G_res, x, y, mix_ratio_init, mu_init, sigma_init)

#Showing the result of spect_em_gmm()
print(cbind(c(mu_init), c(sigma_init), c(mix_ratio_init)))
print(cbind(SP_EM_G_res$mu, SP_EM_G_res$sigma, SP_EM_G_res$mix_ratio))
print(cbind(true_mu, true_sigma, true_mix_ratio))

```

---

show\_lmm\_curve

*Visualization of the result of spect\_em\_lmm*


---

## Description

Visualization of the result of spect\_em\_lmm().

## Usage

```
show_lmm_curve(spect_em_lmm_res, x, y, mix_ratio_init, mu_init, gam_init)
```

## Arguments

spect_em_lmm_res	data set obtained by spect_em_lmm()
x	measurement steps
y	intensity
mix_ratio_init	initial values of the mixture ratio of the components
mu_init	initial values of the mean of the components
gam_init	initial values of the scale parameter of the components

**Details**

Perform a visualization of fitting curve estimated by Lorentz mixture model.

**Value**

Show the fitting curve and variation of the parameters.

**References**

Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2019). Spectrum adapted expectation-maximization algorithm for high-throughput peak shift analysis. *Science and technology of advanced materials*, 20(1), 733-745.

Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2021). Spectrum adapted expectation-conditional maximization algorithm for extending high-throughput peak separation method in XPS analysis. *Science and Technology of Advanced Materials: Methods*, 1(1), 45-55.

**Examples**

```
#generating the synthetic spectral data based on three component Lorentz mixture model.
x          <- seq(0, 100, by = 0.5)
true_mu    <- c(35, 50, 65)
true_gam   <- c(3, 3, 3)
true_mix_ratio <- rep(1/3, 3)
degree     <- 4

#Normalized Lorentz distribution
dCauchy <- function(x, mu, gam) {
  (dcauchy(x, mu, gam)) / sum(dcauchy(x, mu, gam))
}

y <- c(true_mix_ratio[1] * dCauchy(x = x, mu = true_mu[1], gam = true_gam[1])*10^degree +
  true_mix_ratio[2] * dCauchy(x = x, mu = true_mu[2], gam = true_gam[2])*10^degree +
  true_mix_ratio[3] * dCauchy(x = x, mu = true_mu[3], gam = true_gam[3])*10^degree)

plot(y~x, main = "genrated synthetic spectral data")

#Peak fitting by EMpeaksR
#Initial values
K <- 3

mix_ratio_init <- c(0.2, 0.4, 0.4)
mu_init       <- c(20, 40, 70)
gam_init      <- c(2, 5, 4)

#Coducting calculation
SP_ECM_L_res <- spect_em_lmm(x, y, mu = mu_init, gam = gam_init, mix_ratio = mix_ratio_init,
  conv.cri = 1e-2, maxit = 2000)

#Plot fitting curve and trace plot of parameters
show_lmm_curve(SP_ECM_L_res, x, y, mix_ratio_init, mu_init, gam_init)
```

```
#Showing the result of spect_em_lmm()
print(cbind(c(mu_init), c(gam_init), c(mix_ratio_init)))
print(cbind(SP_ECM_L_res$mu, SP_ECM_L_res$gam, SP_ECM_L_res$mix_ratio))
print(cbind(true_mu, true_gam, true_mix_ratio))
```

---

show_pvmm_curve	<i>Visualization of the result of spect_em_pvmm</i>
-----------------	---

---

## Description

Visualization of the result of spect\_em\_pvmm().

## Usage

```
show_pvmm_curve(spect_em_pvmm_res, x, y, mix_ratio_init, mu_init, sigma_init, eta_init)
```

## Arguments

spect_em_pvmm_res	data set obtained by spect_em_pvmm()
x	measurement steps
y	intensity
mix_ratio_init	initial values of the mixture ratio of the components
mu_init	initial values of the mean of the components
sigma_init	initial values of the standard deviation of the components
eta_init	initial values of the mixing ratio of Gauss and Lorentz distribution

## Details

Perform a visualization of fitting curve estimated by Pseudo-Voigt mixture model.

## Value

Show the fitting curve and variation of the parameters.

## References

Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2019). Spectrum adapted expectation-maximization algorithm for high-throughput peak shift analysis. *Science and technology of advanced materials*, 20(1), 733-745.

Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2021). Spectrum adapted expectation-conditional maximization algorithm for extending high-throughput peak separation method in XPS analysis. *Science and Technology of Advanced Materials: Methods*, 1(1), 45-55.



**Examples**

```

#generating the synthetic spectral data based on three component Pseudo-Voigt mixture model.
x          <- seq(0, 100, by = 0.5)
true_mu    <- c(35, 50, 65)
true_sigma <- c(3, 3, 3)
true_eta   <- c(0.3, 0.8, 0.5)
true_mix_ratio <- rep(1/3, 3)
degree     <- 4

#Normalized Pseudo-Voigt distribution
truncated_pv <- function(x, mu, sigma, eta) {
  (eta*dcauchy(x, mu, sqrt(2*log(2))*sigma) + (1-eta)*dnorm(x, mu, sigma)) /
  sum(eta*dcauchy(x, mu, sqrt(2*log(2))*sigma) + (1-eta)*dnorm(x, mu, sigma))
}

y <- c(true_mix_ratio[1]*truncated_pv(x = x,
                                     mu = true_mu[1],
                                     sigma = true_sigma[1],
                                     eta = true_eta[1])*10^degree +
       true_mix_ratio[2]*truncated_pv(x = x,
                                     mu = true_mu[2],
                                     sigma = true_sigma[2],
                                     eta = true_eta[2])*10^degree +
       true_mix_ratio[3]*truncated_pv(x = x,
                                     mu = true_mu[3],
                                     sigma = true_sigma[3],
                                     eta = true_eta[3])*10^degree)

plot(y~x, main = "genrated synthetic spectral data")

#Peak fitting by EMpeaksR
#Initial values
K <- 3

mix_ratio_init <- c(0.2, 0.4, 0.4)
mu_init        <- c(20, 40, 70)
sigma_init     <- c(2, 5, 4)
eta_init       <- c(0.5, 0.4, 0.3)

#Coducting calculation
SP_ECM_PV_res <- spect_em_pvmm(x = x,
                               y = y,
                               mu = mu_init,
                               sigma = sigma_init,
                               eta = eta_init,
                               mix_ratio = mix_ratio_init,
                               conv.cri = 1e-2,
                               maxit = 2000)

#Plot fitting curve and trace plot of parameters
show_pvmm_curve(SP_ECM_PV_res, x, y, mix_ratio_init, mu_init, sigma_init, eta_init)

```

```
#Showing the result of spect_em_pvmm()
print(cbind(c(mu_init), c(sigma_init), c(eta_init), c(mix_ratio_init)))
print(cbind(SP_ECM_PV_res$mu, SP_ECM_PV_res$sigma, SP_ECM_PV_res$eta, SP_ECM_PV_res$mix_ratio))
print(cbind(true_mu, true_sigma, true_eta, true_mix_ratio))
```

---

show\_pvmm\_lback\_curve *Visualization of the result of spect\_em\_pvmm\_lback*

---

## Description

Visualization of the result of spect\_em\_pvmm\_lback().

## Usage

```
show_pvmm_lback_curve(spect_em_pvmm_lback_res,
                      x, y,
                      mix_ratio_init,
                      mu_init,
                      sigma_init,
                      eta_init,
                      x_lower,
                      x_upper)
```

## Arguments

spect_em_pvmm_lback_res	data set obtained by spect_em_pvmm_lback()
x	measurement steps
y	intensity
mu_init	initial values of the mean of the components
sigma_init	initial values of the standard deviation of the components
eta_init	initial values of the mixing ratio of Gauss and Lorentz distribution
mix_ratio_init	initial values of the mixture ratio of the components
x_lower	lower limit of the measurement steps. Default is a minimum of x
x_upper	upper limit of the measurement steps. Default is a maximum of x

## Details

Perform a visualization of fitting curve estimated by pseudo-Voigt mixture model with a linear background.

## Value

Show the fitting curve and variation of the parameters.

## References

Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2019). Spectrum adapted expectation-maximization algorithm for high-throughput peak shift analysis. *Science and technology of advanced materials*, 20(1), 733-745.

Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2021). Spectrum adapted expectation-conditional maximization algorithm for extending high-throughput peak separation method in XPS analysis. *Science and Technology of Advanced Materials: Methods*, 1(1), 45-55.

Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2023). High-throughput XPS spectrum modeling with autonomous background subtraction for 3 d 5/2 peak mapping of SnS. *Science and Technology of Advanced Materials: Methods*, 3(1), 2159753.

## Examples

```
#generating the synthetic spectral data based on three component Pseudo-Voigt mixture model.
x          <- seq(0, 100, by = 0.5)
K          <- 3
true_mu    <- c(35, 50, 65)
true_sigma <- c(3, 3, 3)
true_mix_ratio <- c(0.5/3, 0.5/3, 0.5/3, 0.5)
true_eta   <- c(0.4, 0.6, 0.1)
degree    <- 4

#Normalized Pseudo-Voigt distribution
truncated_pv <- function(x, mu, sigma, eta) {
  (eta*dcauchy(x, mu, sqrt(2*log(2))*sigma) + (1-eta)*dnorm(x, mu, sigma)) /
  sum(eta*dcauchy(x, mu, sqrt(2*log(2))*sigma) + (1-eta)*dnorm(x, mu, sigma))
}

y <- c(true_mix_ratio[1]*truncated_pv(x = x,
                                     mu = true_mu[1],
                                     sigma = true_sigma[1],
                                     eta = true_eta[1])*10^degree +
      true_mix_ratio[2]*truncated_pv(x = x,
                                     mu = true_mu[2],
                                     sigma = true_sigma[2],
                                     eta = true_eta[2])*10^degree +
      true_mix_ratio[3]*truncated_pv(x = x,
                                     mu = true_mu[3],
                                     sigma = true_sigma[3],
                                     eta = true_eta[3])*10^degree +
      true_mix_ratio[4]*(c(500*x + 15000) / sum(500*x + 15000))*10^degree)

plot(y~x, main = "genrated synthetic spectral data")

#Peak fitting by EMpeaksR
#Initial values
mu_init    <- c(30, 40, 60)
sigma_init <- c(4, 4, 4)
mix_ratio_init <- rep(1/(length(mu_init)+3), length(mu_init)+3)
eta_init   <- c(1, 1, 1)
```

```

#Conducting calculation
SP_ECM_PV_LBACK_res <- spect_em_pvmm_lback(x = x,
                                           y = y,
                                           mu = mu_init,
                                           sigma = sigma_init,
                                           eta = eta_init,
                                           mix_ratio = mix_ratio_init,
                                           x_lower = min(x),
                                           x_upper = max(x),
                                           conv.cri = 1e-2,
                                           maxit = 2000)

#Plot fitting curve and trace plot of parameters
show_pvmm_lback_curve(spect_em_pvmm_lback_res = SP_ECM_PV_LBACK_res,
                      x = x,
                      y = y,
                      mix_ratio_init = mix_ratio_init,
                      mu_init = mu_init,
                      sigma_init = sigma_init,
                      eta_init = eta_init,
                      x_lower = min(x),
                      x_upper = max(x))

#Showing the result of spect_em_pvmm_lback()
print(cbind(SP_ECM_PV_LBACK_res$mu, SP_ECM_PV_LBACK_res$sigma, SP_ECM_PV_LBACK_res$eta,
            SP_ECM_PV_LBACK_res$mix_ratio[1:K]))

print(cbind(true_mu, true_sigma, true_eta, true_mix_ratio[1:K]))

```

---

spect\_em\_dsgmm

*Spectrum adapted ECM algorithm by DSGMM*

---

### Description

Perform a peak fitting based on the spectrum adapted ECM algorithm by Doniach-Sunjic-Gauss mixture model.

### Usage

```
spect_em_dsgmm(x, y, mu, sigma, alpha, eta, mix_ratio, conv.cri, maxit)
```

### Arguments

x	measurement steps
y	intensity
mu	mean of the components
sigma	standard deviation of the components

alpha	asymmetric parameter of the component
eta	mixing ratio of Gauss and Lorentz distribution
mix_ratio	mixture ratio of the components
conv.cri	criterion of the convergence
maxit	maximum number of the iteration

### Details

Peak fitting is conducted by spectrum adapted ECM algorithm.

### Value

mu	estimated mean of the components
sigma	estimated standard deviation of the components
alpha	estimated asymmetric parameter of the components
eta	estimated mixing ratio of Gauss and Lorentz distribution
mix_ratio	estimated mixture ratio of the components
it	number of the iteration to reach the convergence
LL	variation of the weighted log likelihood values
MU	variation of mu
SIGMA	variation of sigma
ALPHA	variation of alpha
ETA	variation of beta
MIX_RATIO	variation of mix_ratio
W_K	decomposed component of the spectral data
convergence	message for the convergence in the calculation
cal_time	calculation time to complete the peak fitting. Unit is seconds

### References

Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2019). Spectrum adapted expectation-maximization algorithm for high-throughput peak shift analysis. *Science and technology of advanced materials*, 20(1), 733-745.

Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2021). Spectrum adapted expectation-conditional maximization algorithm for extending high-throughput peak separation method in XPS analysis. *Science and Technology of Advanced Materials: Methods*, 1(1), 45-55.

**Examples**

```

#generating the synthetic spectral data based on three component Doniach-Sunjic-Gauss mixture model.
x          <- seq(0, 100, by = 0.5)
true_mu    <- c(20, 50, 80)
true_sigma <- c(3, 3, 3)
true_alpha <- c(0.1, 0.3, 0.1)
true_eta   <- c(0.4, 0.6, 0.1)
true_mix_ratio <- rep(1/3, 3)
degree     <- 4

#truncated Doniach-Sunjic-Gauss
truncated_dsg <- function(x, mu, sigma, alpha, eta) {
  ((eta*(((gamma(1-alpha)) /
  ((x-mu)^2+(sqrt(2*log(2))*sigma)^2)^((1-alpha)/2)) *
  cos((pi*alpha/2)+(1-alpha)*atan((x-mu) /
  (sqrt(2*log(2))*sigma)))) + (1-eta)*dnorm(x, mu, sigma)) /
  sum( ((eta*(((gamma(1-alpha)) /
  ((x-mu)^2+(sqrt(2*log(2))*sigma)^2)^((1-alpha)/2)) *
  cos((pi*alpha/2)+(1-alpha)*atan((x-mu) /
  (sqrt(2*log(2))*sigma)))) + (1-eta)*dnorm(x, mu, sigma)))
}

y <- c(true_mix_ratio[1]*truncated_dsg(x = x,
                                     mu = true_mu[1],
                                     sigma = true_sigma[1],
                                     alpha = true_alpha[1],
                                     eta = true_eta[1])*10^degree +
      true_mix_ratio[2]*truncated_dsg(x = x,
                                     mu = true_mu[2],
                                     sigma = true_sigma[2],
                                     alpha = true_alpha[2],
                                     eta = true_eta[2])*10^degree +
      true_mix_ratio[3]*truncated_dsg(x = x,
                                     mu = true_mu[3],
                                     sigma = true_sigma[3],
                                     alpha = true_alpha[3],
                                     eta = true_eta[3])*10^degree)

plot(y~x, main = "genrated synthetic spectral data")

#Peak fitting by EMpeaksR
#Initial values
K <- 3
mix_ratio_init <- c(0.2, 0.4, 0.4)
mu_init       <- c(20, 40, 70)
sigma_init    <- c(4, 3, 2)
alpha_init    <- c(0.3, 0.2, 0.4)
eta_init      <- c(0.5, 0.4, 0.3)

#Coducting calculation
SP_ECM_DSG_res <- spect_em_dsgmm(x = x,
                                 y = y,

```

```

                                mu = mu_init,
                                sigma = sigma_init,
                                alpha = alpha_init,
                                eta = eta_init,
                                mix_ratio = mix_ratio_init,
                                conv.cri = 1e-2,
                                maxit = 2000)

#Plot fitting curve and trace plot of parameters
show_dsgmm_curve(SP_ECM_DSG_res,
                 x,
                 y,
                 mix_ratio_init,
                 mu_init,
                 sigma_init,
                 alpha_init,
                 eta_init)

#Showing the result of spect_em_dsgmm()
print(cbind(c(mu_init),
            c(sigma_init),
            c(alpha_init),
            c(eta_init),
            c(mix_ratio_init)))

print(cbind(SP_ECM_DSG_res$mu,
            SP_ECM_DSG_res$sigma,
            SP_ECM_DSG_res$alpha,
            SP_ECM_DSG_res$eta,
            SP_ECM_DSG_res$mix_ratio))

print(cbind(true_mu,
            true_sigma,
            true_alpha,
            true_eta,
            true_mix_ratio))

```

---

spect\_em\_gmm

*Spectrum adapted EM algorithm by GMM*


---

### Description

Perform a peak fitting based on the spectrum adapted EM algorithm by Gaussian mixture model.

### Usage

```
spect_em_gmm(x, y, mu, sigma, mix_ratio, conv.cri, maxit)
```

**Arguments**

x	measurement steps
y	intensity
mu	mean of the components
sigma	standard deviation of the components
mix_ratio	mixture ratio of the components
conv.cri	criterion of the convergence
maxit	maximum number of the iteration

**Details**

Peak fitting is conducted by spectrum adapted EM algorithm.

**Value**

mu	estimated mean of the components
sigma	estimated standard deviation of the components
mix_ratio	estimated mixture ratio of the components
it	number of the iteration to reach the convergence
LL	variation of the weighted log likelihood values
MU	variation of mu
SIGMA	variation of sigma
MIX_RATIO	variation of mix_ratio
W_K	decomposed component of the spectral data
convergence	message for the convergence in the calculation
cal_time	calculation time to complete the peak fitting. Unit is seconds

**References**

Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2019). Spectrum adapted expectation-maximization algorithm for high-throughput peak shift analysis. *Science and technology of advanced materials*, 20(1), 733-745.

**Examples**

```
#generating the synthetic spectral data based on three component Gaussian mixture model.
x <- seq(0, 100, by = 0.5)
true_mu <- c(35, 50, 65)
true_sigma <- c(3, 3, 3)
true_mix_ratio <- rep(1/3, 3)
degree <- 4

y <- c(true_mix_ratio[1] * dnorm(x = x, mean = true_mu[1], sd = true_sigma[1])*10^degree +
      true_mix_ratio[2] * dnorm(x = x, mean = true_mu[2], sd = true_sigma[2])*10^degree +
      true_mix_ratio[3] * dnorm(x = x, mean = true_mu[3], sd = true_sigma[3])*10^degree)
```



```

plot(y~x, main = "genrated synthetic spectral data")

#Peak fitting by EMpeaksR
#Initial values
K <- 3

mix_ratio_init <- c(0.2, 0.4, 0.4)
mu_init        <- c(20, 40, 70)
sigma_init     <- c(2, 5, 4)

#Coducting calculation
SP_EM_G_res <- spect_em_gmm(x, y, mu = mu_init, sigma = sigma_init, mix_ratio = mix_ratio_init,
                           conv.cri = 1e-2, maxit = 2000)

#Plot fitting curve and trace plot of parameters
show_gmm_curve(SP_EM_G_res, x, y, mix_ratio_init, mu_init, sigma_init)

#Showing the result of spect_em_gmm()
print(cbind(c(mu_init), c(sigma_init), c(mix_ratio_init)))
print(cbind(SP_EM_G_res$mu, SP_EM_G_res$sigma, SP_EM_G_res$mix_ratio))
print(cbind(true_mu, true_sigma, true_mix_ratio))

```

---

spect\_em\_lmm

*Spectrum adapted ECM algorithm by LMM*


---

## Description

Perform a peak fitting based on the spectrum adapted ECM algorithm by Lorentz mixture model.

## Usage

```
spect_em_lmm(x, y, mu, gam, mix_ratio, conv.cri, maxit)
```

## Arguments

x	measurement steps
y	intensity
mu	mean of the components
gam	scale parameter of the components
mix_ratio	mixture ratio of the components
conv.cri	criterion of the convergence
maxit	maximum number of the iteration

## Details

Peak fitting is conducted by spectrum adapted ECM algorithm.

**Value**

mu	estimated mean of the components
gam	estimated scale parameter of the components
mix_ratio	estimated mixture ratio of the components
it	number of the iteration to reach the convergence
LL	variation of the weighted log likelihood values
MU	variation of mu
GAM	variation of gam
MIX_RATIO	variation of mix_ratio
W_K	decomposed component of the spectral data
convergence	message for the convergence in the calculation
cal_time	calculation time to complete the peak fitting. Unit is seconds

**References**

Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2019). Spectrum adapted expectation-maximization algorithm for high-throughput peak shift analysis. *Science and technology of advanced materials*, 20(1), 733-745.

Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2021). Spectrum adapted expectation-conditional maximization algorithm for extending high-throughput peak separation method in XPS analysis. *Science and Technology of Advanced Materials: Methods*, 1(1), 45-55.

**Examples**

```
#generating the synthetic spectral data based on three component Lorentz mixture model.
x      <- seq(0, 100, by = 0.5)
true_mu <- c(35, 50, 65)
true_gam <- c(3, 3, 3)
true_mix_ratio <- rep(1/3, 3)
degree <- 4

#Normalized Lorentz distribution
dCauchy <- function(x, mu, gam) {
  (dcauchy(x, mu, gam)) / sum(dcauchy(x, mu, gam))
}

y <- c(true_mix_ratio[1] * dCauchy(x = x, mu = true_mu[1], gam = true_gam[1])*10^degree +
  true_mix_ratio[2] * dCauchy(x = x, mu = true_mu[2], gam = true_gam[2])*10^degree +
  true_mix_ratio[3] * dCauchy(x = x, mu = true_mu[3], gam = true_gam[3])*10^degree)

plot(y~x, main = "genrated synthetic spectral data")

#Peak fitting by EMpeaksR
#Initial values
K <- 3

mix_ratio_init <- c(0.2, 0.4, 0.4)
```

```

mu_init      <- c(20, 40, 70)
gam_init     <- c(2, 5, 4)

#Conducting calculation
SP_ECM_L_res <- spect_em_lmm(x, y, mu = mu_init, gam = gam_init, mix_ratio = mix_ratio_init,
                             conv.cri = 1e-2, maxit = 2000)

#Plot fitting curve and trace plot of parameters
show_lmm_curve(SP_ECM_L_res, x, y, mix_ratio_init, mu_init, gam_init)

#Showing the result of spect_em_lmm()
print(cbind(c(mu_init), c(gam_init), c(mix_ratio_init)))
print(cbind(SP_ECM_L_res$mu, SP_ECM_L_res$gam, SP_ECM_L_res$mix_ratio))
print(cbind(true_mu, true_gam, true_mix_ratio))

```

---

spect\_em\_pvmm

*Spectrum adapted ECM algorithm by PVMM*


---

### Description

Perform a peak fitting based on the spectrum adapted ECM algorithm by Pseudo-Voigt mixture model.

### Usage

```

spect_em_pvmm(x, y, mu, sigma, eta, mix_ratio, conv.cri, maxit)

```

### Arguments

x	measurement steps
y	intensity
mu	mean of the components
sigma	standard deviation of the components
eta	mixing ratio of Gauss and Lorentz distribution
mix_ratio	mixture ratio of the components
conv.cri	criterion of the convergence
maxit	maximum number of the iteration

### Details

Peak fitting is conducted by spectrum adapted ECM algorithm.

**Value**

mu	estimated mean of the components
sigma	estimated standard deviation of the components
eta	estimated mixing ratio of Gauss and Lorentz distribution
mix_ratio	estimated mixture ratio of the components
it	number of the iteration to reach the convergence
LL	variation of the weighted log likelihood values
MU	variation of mu
SIGMA	variation of sigma
ETA	variation of beta
MIX_RATIO	variation of mix_ratio
W_K	decomposed component of the spectral data
convergence	message for the convergence in the calculation
cal_time	calculation time to complete the peak fitting. Unit is seconds

**References**

Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2019). Spectrum adapted expectation-maximization algorithm for high-throughput peak shift analysis. *Science and technology of advanced materials*, 20(1), 733-745.

Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2021). Spectrum adapted expectation-conditional maximization algorithm for extending high-throughput peak separation method in XPS analysis. *Science and Technology of Advanced Materials: Methods*, 1(1), 45-55.

**Examples**

```
#generating the synthetic spectral data based on three component Pseudo-Voigt mixture model.
x <- seq(0, 100, by = 0.5)
true_mu <- c(35, 50, 65)
true_sigma <- c(3, 3, 3)
true_eta <- c(0.3, 0.8, 0.5)
true_mix_ratio <- rep(1/3, 3)
degree <- 4

#Normalized Pseudo-Voigt distribution
truncated_pv <- function(x, mu, sigma, eta) {
  (eta*dcauchy(x, mu, sqrt(2*log(2))*sigma) + (1-eta)*dnorm(x, mu, sigma)) /
  sum(eta*dcauchy(x, mu, sqrt(2*log(2))*sigma) + (1-eta)*dnorm(x, mu, sigma))
}

y <- c(true_mix_ratio[1]*truncated_pv(x = x,
                                     mu = true_mu[1],
                                     sigma = true_sigma[1],
                                     eta = true_eta[1])*10^degree +
      true_mix_ratio[2]*truncated_pv(x = x,
                                     mu = true_mu[2],
```

```

                                sigma = true_sigma[2],
                                eta = true_eta[2])*10^degree +
true_mix_ratio[3]*truncated_pv(x = x,
                                mu = true_mu[3],
                                sigma = true_sigma[3],
                                eta = true_eta[3])*10^degree)

plot(y~x, main = "genrated synthetic spectral data")

#Peak fitting by EMpeaksR
#Initial values
K <- 3

mix_ratio_init <- c(0.2, 0.4, 0.4)
mu_init        <- c(20, 40, 70)
sigma_init     <- c(2, 5, 4)
eta_init       <- c(0.5, 0.4, 0.3)

#Conducting calculation
SP_ECM_PV_res <- spect_em_pvmm(x = x,
                               y = y,
                               mu = mu_init,
                               sigma = sigma_init,
                               eta = eta_init,
                               mix_ratio = mix_ratio_init,
                               conv.cri = 1e-2,
                               maxit = 2000)

#Plot fitting curve and trace plot of parameters
show_pvmm_curve(SP_ECM_PV_res, x, y, mix_ratio_init, mu_init, sigma_init, eta_init)

#Showing the result of spect_em_pvmm()
print(cbind(c(mu_init), c(sigma_init), c(eta_init), c(mix_ratio_init)))
print(cbind(SP_ECM_PV_res$mu, SP_ECM_PV_res$sigma, SP_ECM_PV_res$eta, SP_ECM_PV_res$mix_ratio))
print(cbind(true_mu, true_sigma, true_eta, true_mix_ratio))

```

---

spect\_em\_pvmm\_lback     *Spectrum adapted ECM algorithm by PVMM with a linear background*

---

### Description

Perform a peak fitting based on the spectrum adapted ECM algorithm by pseudo-Voigt mixture model with a linear background.

### Usage

```
spect_em_pvmm_lback(x, y, mu, sigma, eta, mix_ratio, x_lower, x_upper, conv.cri, maxit)
```

**Arguments**

x	measurement steps
y	intensity
mu	mean of the components
sigma	standard deviation of the components
eta	mixing ratio of Gauss and Lorentz distribution
mix_ratio	mixture ratio of the components
x_lower	lower limit of the measurement steps. Default is a minimum of x
x_upper	upper limit of the measurement steps. Default is a maximum of x
conv.cri	criterion of the convergence
maxit	maximum number of the iteration

**Details**

Peak fitting is conducted by spectrum adapted ECM algorithm.

**Value**

mu	estimated mean of the components
sigma	estimated standard deviation of the components
eta	estimated mixing ratio of Gauss and Lorentz distribution
mix_ratio	estimated mixture ratio of the components
it	number of the iteration to reach the convergence
LL	variation of the weighted log likelihood values
MU	variation of mu
SIGMA	variation of sigma
ETA	variation of beta
MIX_RATIO	variation of mix_ratio
W_K	decomposed component of the spectral data
convergence	message for the convergence in the calculation
cal_time	calculation time to complete the peak fitting. Unit is seconds

**References**

- Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2019). Spectrum adapted expectation-maximization algorithm for high-throughput peak shift analysis. *Science and technology of advanced materials*, 20(1), 733-745.
- Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2021). Spectrum adapted expectation-conditional maximization algorithm for extending high-throughput peak separation method in XPS analysis. *Science and Technology of Advanced Materials: Methods*, 1(1), 45-55.
- Matsumura, T., Nagamura, N., Akaho, S., Nagata, K., & Ando, Y. (2023). High-throughput XPS spectrum modeling with autonomous background subtraction for 3 d 5/2 peak mapping of SnS. *Science and Technology of Advanced Materials: Methods*, 3(1), 2159753.

**Examples**

```

#generating the synthetic spectral data based on three component Pseudo-Voigt mixture model.
x          <- seq(0, 100, by = 0.5)
K          <- 3
true_mu    <- c(35, 50, 65)
true_sigma <- c(3, 3, 3)
true_mix_ratio <- c(0.5/3, 0.5/3, 0.5/3, 0.5)
true_eta   <- c(0.4, 0.6, 0.1)
degree    <- 4

#Normalized Pseudo-Voigt distribution
truncated_pv <- function(x, mu, sigma, eta) {
  (eta*dcauchy(x, mu, sqrt(2*log(2))*sigma) + (1-eta)*dnorm(x, mu, sigma)) /
  sum(eta*dcauchy(x, mu, sqrt(2*log(2))*sigma) + (1-eta)*dnorm(x, mu, sigma))
}

y <- c(true_mix_ratio[1]*truncated_pv(x = x,
                                     mu = true_mu[1],
                                     sigma = true_sigma[1],
                                     eta = true_eta[1])*10^degree +
        true_mix_ratio[2]*truncated_pv(x = x,
                                     mu = true_mu[2],
                                     sigma = true_sigma[2],
                                     eta = true_eta[2])*10^degree +
        true_mix_ratio[3]*truncated_pv(x = x,
                                     mu = true_mu[3],
                                     sigma = true_sigma[3],
                                     eta = true_eta[3])*10^degree +
        true_mix_ratio[4]*(c(500*x + 15000) / sum(500*x + 15000))*10^degree)

plot(y~x, main = "generated synthetic spectral data")

#Peak fitting by EMpeaksR
#Initial values
mu_init    <- c(30, 40, 60)
sigma_init <- c(4, 4, 4)
mix_ratio_init <- rep(1/(length(mu_init)+3), length(mu_init)+3)
eta_init   <- c(1, 1, 1)

#Conducting calculation
SP_ECM_PV_LBACK_res <- spect_em_pvmm_lback(x = x,
                                             y = y,
                                             mu = mu_init,
                                             sigma = sigma_init,
                                             eta = eta_init,
                                             mix_ratio = mix_ratio_init,
                                             x_lower = min(x),
                                             x_upper = max(x),
                                             conv.cri = 1e-2,
                                             maxit = 2000)

#Plot fitting curve and trace plot of parameters

```

```
show_pvmm_lback_curve(spect_em_pvmm_lback_res = SP_ECM_PV_LBACK_res,  
                      x = x,  
                      y = y,  
                      mix_ratio_init = mix_ratio_init,  
                      mu_init = mu_init,  
                      sigma_init = sigma_init,  
                      eta_init = eta_init,  
                      x_lower = min(x),  
                      x_upper = max(x))  
  
#Showing the result of spect_em_pvmm_lback()  
print(cbind(SP_ECM_PV_LBACK_res$mu, SP_ECM_PV_LBACK_res$sigma, SP_ECM_PV_LBACK_res$eta,  
           SP_ECM_PV_LBACK_res$mix_ratio[1:K]))  
  
print(cbind(true_mu, true_sigma, true_eta, true_mix_ratio[1:K]))
```



# Index

[show\\_dsgmm\\_curve](#), [2](#)  
[show\\_gmm\\_curve](#), [5](#)  
[show\\_lmm\\_curve](#), [6](#)  
[show\\_pvmm\\_curve](#), [8](#)  
[show\\_pvmm\\_lback\\_curve](#), [10](#)  
[spect\\_em\\_dsgmm](#), [12](#)  
[spect\\_em\\_gmm](#), [15](#)  
[spect\\_em\\_lmm](#), [17](#)  
[spect\\_em\\_pvmm](#), [19](#)  
[spect\\_em\\_pvmm\\_lback](#), [21](#)